

# 10 | Cofibrations

**10.1 Definition.** A map  $i: A \rightarrow X$  has the *homotopy extension property* for a space  $Y$  if for any commutative diagram of the form

$$\begin{array}{ccc} Y & \xleftarrow{\bar{f}} & X \\ \uparrow \text{ev}_0 & \nearrow \bar{h} & \uparrow i \\ PY & \xleftarrow{h} & A \end{array}$$

there exists a map  $\bar{h}: X \times [0, 1] \rightarrow E$  such that  $\bar{h}i = \bar{f}$  and  $p\bar{h} = h$ . Here  $PY$  is the path space of  $Y$  and  $\text{ev}_0: PY \rightarrow Y$  is the evaluation at 0 map:  $\text{ev}_0(\omega) = \omega(0)$ .

Equivalently,  $i: A \rightarrow X$  has the homotopy extension property for  $Y$  if given any map  $\bar{f}: X \rightarrow Y$  and a homotopy  $h^\sharp: A \times [0, 1] \rightarrow Y$  such that  $h_0^\sharp = \bar{f}i$  we can find a homotopy  $\bar{h}^\sharp: X \times [0, 1]$ , such that  $\bar{h}_0^\sharp = \bar{f}$  and  $\bar{h}^\sharp(i(a), t) = h^\sharp(a, t)$  for all  $(a, t) \in A \times [0, 1]$ .

In this setting we will say that  $\bar{h}^\sharp$  is an extension of  $h^\sharp$  beginning at  $\bar{f}$ .

**10.2 Definition.** A map  $i: A \rightarrow X$  is a *cofibration* if it has the homotopy extension property for any space  $Y$ . In such case we also say that the space  $X/i(A)$  is the *cofiber* of  $i$ .

**10.3 Example.** By Theorem 2.14 if  $(X, A)$  is a relative CW complex then the inclusion  $i: A \hookrightarrow X$  is a cofibration.

Recall that the mapping cylinder of a map  $f: X \rightarrow Y$  is the quotient space

$$M_f = (X \times [0, 1] \sqcup Y) / \sim$$

where  $(x, 0) \sim f(x)$  for all  $x \in X$ . We have a map  $s_f: M_f \rightarrow Y \times [0, 1]$  such that  $s_f(x, t) = (f(x), t)$  for  $(x, t) \in X \times [0, 1]$  and  $s_f(y) = (y, 0)$  for  $y \in Y$ .

**10.4 Proposition.** For a map  $i: A \rightarrow X$  the following conditions are equivalent:

- 1) The map  $i$  is a cofibration.
- 2) The map  $i$  has the homotopy extension property for the space  $M_i$
- 3) There exists a map  $r_f: X \times [0, 1] \rightarrow M_i$  such that  $r_f s_f = \text{id}_{M_i}$

*Proof.* Exercise. □

**10.5 Corollary.** If  $i: A \rightarrow X$  is a cofibration then  $i$  is an embedding.

*Proof.* Exercise. Use condition 3) in Proposition 10.4. □

**10.6 Proposition.** Given any map  $f: X \rightarrow Y$  the map  $i_f: X \rightarrow M_f$  given by  $i_f(x) = (x, 1)$  is a cofibration.

*Proof.* Exercise. □

**10.7 Note.** Given a map  $f: X \rightarrow Y$ , let  $d_f: M_f \rightarrow Y$  be the strong deformation retraction. As a consequence of Proposition 10.6, we have a commutative diagram

$$\begin{array}{ccc} Y & \xleftarrow[d_f]{\simeq} & M_f \\ & \nwarrow f \quad \nearrow i_f & \\ & X & \end{array}$$

where  $i_f$  is a cofibration. A homotopy inverse of  $d_f$  is given by the inclusion map  $j_f: Y \rightarrow M_f$ .

**10.8 Note.** Recall that the mapping cone of a map  $f: X \rightarrow Y$  is the space  $C_f = M_f/X \times \{1\}$ . The space  $C_f$  is the cofiber of the cofibration  $i_f: X \rightarrow M_f$ .

**10.9 Coexact Puppe sequence.** The construction of the coexact Puppe sequence of a map is dual to the construction of the exact Puppe sequence given in Chapter 9.

As in Chapter 9 we will be interested here in pointed spaces and homotopy classes of maps that preserve basepoints. In this case we will use a slightly weakened version of a cofibration: a map of pointed spaces  $i: (A, a_0) \rightarrow (X, x_0)$  is a cofibration if has the homotopy extension property for all pointed maps  $(X, x_0) \rightarrow (Y, y_0)$  and pointed homotopies  $A \times [0, 1] \rightarrow Y$ . In this context we modify the constructions of the mapping mapping cylinder and the mapping cone as follows:

**10.10 Definition.** For a map of pointed spaces  $f: (X, x_0) \rightarrow (Y, y_0)$  the *reduced mapping cylinder* of  $f$  is the space  $\bar{M}_f = M_f/\{x_0\} \times [0, 1]$ . The *reduced mapping cone* is the space  $\bar{C}_f = \bar{M}_f/X \times \{1\}$ .

The reduced mapping cylinder and mapping cone come with a natural choice of basepoints. As in (10.7) for any map  $f: (X, x_0) \rightarrow (Y, y_0)$  we have a commutative diagram

$$\begin{array}{ccc} Y & \xleftarrow[d_f]{\simeq} & \bar{M}_f \\ & \nearrow f & \nwarrow i_f \\ & X & \end{array}$$

where  $i_f$  is a pointed cofibration and  $d_f$  is a pointed homotopy equivalence. Also,  $\bar{C}_f$  is the cofiber of  $i_f$ .

**10.11 Definition.** A sequence of maps of spaces

$$(X_0, x_0) \xrightarrow{f_0} (X_1, x_1) \xrightarrow{f_1} (X_2, x_2)$$

is *coexact at  $X_1$*  is for any pointed space  $(Y, y_0)$  the sequence pointed sets

$$[X_2, Y]_* \xrightarrow{f_1^*} [X_1, Y]_* \xrightarrow{f_0^*} [X_0, Y]_*$$

is exact at  $[X_1, Y]_*$ .

**10.12 Proposition.** If  $i: A \rightarrow X$  is a cofibration,  $q: X \rightarrow X/i(A)$  is the quotient map,  $x_0 \in A$  then the sequence  $(A, x_0) \xrightarrow{i} (X, i(x_0)) \xrightarrow{q} (X/A, qi(x_0))$  is coexact at  $X$ .

For any map  $f: (X, x_0) \rightarrow (Y, y_0)$  consider the sequence

$$X \xrightarrow{f} Y \xrightarrow{q(f)} \bar{C}_f$$

where  $q(f)(y) = (y, 0)$ . Since this sequence is homotopy equivalent to the cofibration sequence  $X \xrightarrow{i_f} \bar{M}_f \rightarrow \bar{C}_f$ , it is coexact at  $Y$ . Continuing this construction inductively we obtain a coexact sequence

$$X \xrightarrow{f} Y \xrightarrow{q(f)} \bar{C}_f \xrightarrow{q^2(f)} \bar{C}_{q(f)} \xrightarrow{q^3(f)} \bar{C}_{q^2(f)} \xrightarrow{q^4(f)} \bar{C}_{q^3(f)} \rightarrow \dots \quad (*)$$

As in Chapter 9 our goal will be to show that this sequence admits a more convenient description. This will depend on two facts that dualize Proposition 9.4 and Corollary 8.18

**10.13 Proposition.** For any map  $f: (X, x_0) \rightarrow (Y, y_0)$  the map  $q(f): X \rightarrow \bar{C}_f$  is a cofibration.

*Proof.* Exercise. □

**10.14 Proposition.** If  $f: (X, x_0) \rightarrow (Y, y_0)$  is a cofibration then the quotient map

$$\bar{C}_f \rightarrow Y/f(X)$$

is a homotopy equivalence.

*Proof.* Exercise. □

Notice that  $\bar{C}_f/q(f) \cong \Sigma X$ , where  $\Sigma X$  is the reduced suspension of  $X$ . In this way we obtain:

**10.15 Proposition.** *For any map of pointed spaces  $f: (X, x_0) \rightarrow (Y, y_0)$  we have a commutative diagram*

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{q(f)} & \bar{C}_f & \xrightarrow{q^2(f)} & \bar{C}_{q(f)} \\ & & & & \searrow g & & \downarrow \simeq \\ & & & & & & \Sigma X \end{array}$$

Applying Proposition 10.14 iteratively to the sequence  $(*)$  we get homotopy equivalences

$$\begin{aligned} \bar{C}_{q(f)} &\xrightarrow{\simeq} \Sigma X \\ \bar{C}_{q^2(f)} &\xrightarrow{\simeq} \Sigma Y \\ \bar{C}_{q^3(f)} &\xrightarrow{\simeq} \Sigma \bar{C}_f \\ \bar{C}_{q^4(f)} &\xrightarrow{\simeq} \Sigma \bar{C}_{q(f)} \simeq \Sigma^2 X \\ \bar{C}_{q^5(f)} &\xrightarrow{\simeq} \Sigma \bar{C}_{q^2(f)} \simeq \Sigma^2 Y \\ &\dots \quad \dots \quad \dots \quad \dots \quad \dots \end{aligned}$$

Moreover, one can check that the following diagram commutes up to homotopy:

$$\begin{array}{ccccccccccccccc} X & \xrightarrow{f} & Y & \xrightarrow{q(f)} & \bar{C}_f & \xrightarrow{q^2(f)} & \bar{C}_{q(f)} & \xrightarrow{q^3(f)} & \bar{C}_{q^2(f)} & \xrightarrow{q^4(f)} & \bar{C}_{q^3(f)} & \xrightarrow{q^5(f)} & \bar{C}_{q^4(f)} & \longrightarrow & \dots \\ \parallel & & \parallel & & \parallel & & \wr & & \wr & & \wr & & \wr & & \\ X & \xrightarrow{f} & Y & \xrightarrow{q(f)} & \bar{C}_f & \xrightarrow{g} & \Sigma X & \xrightarrow{\Sigma f} & \Sigma Y & \xrightarrow{\Sigma q(f)} & \Sigma \bar{C}_f & \xrightarrow{\Sigma g} & \Sigma^2 X & \longrightarrow & \dots \end{array} \quad (**)$$

**10.16 Definition.** The sequence in the lower row of the diagram  $(**)$  is called the *Puppe coexact sequence* associated to the map  $f$ .

As a consequence, for any map of pointed spaces  $f: (X, x_0) \rightarrow (Y, y_0)$  and any pointed space  $(Z, z_0)$  we obtain a long exact sequence of sets:

$$[X, Z]_* \xleftarrow{f^*} [Y, Z]_* \xleftarrow{q(f)^*} [\bar{C}_f, Z]_* \xleftarrow{g^*} [\Sigma X, Z]_* \xleftarrow{\Sigma f^*} [\Sigma Y, Z]_* \xleftarrow{\Sigma q(f)^*} [\Sigma \bar{C}_f, Z]_* \xleftarrow{\Sigma g^*} [\Sigma^2 X, Z]_* \longleftarrow \dots \quad (\star)$$

Starting with  $[\Sigma X, Z]_*$  the sets in this sequence have a group structure defined by the suspension, and all maps are homomorphisms of groups. Starting with  $[\Sigma^2, Z]_*$  all groups are abelian.

**10.17 Note.** 1) Using the adjunction  $\text{adj}: [\Sigma X, Y]_* \xrightarrow{\cong} [X, \Omega Y]_*$  as in (9.15) we can rewrite the sequence (X) in the form

$$[X, Z]_* \xleftarrow{f^*} [Y, Z]_* \xleftarrow{q(f)^*} [\bar{C}_f, Z]_* \xleftarrow{g^*} [X, \Omega Z]_* \xleftarrow{f^*} [Y, \Omega Z]_* \xleftarrow{q(f)^*} [\bar{C}_f, \Omega Z]_* \xleftarrow{g^*} [X, \Omega^2 Z]_* \leftarrow \dots$$

In this setting, groups structures are induced the multiplication in loop spaces.

2) Assume that the map  $f: (X, x_0) \rightarrow (Y, y_0)$  is a cofibration. Using Corollary 10.5 we can then assume that  $X$  is a subspace of  $Y$  and that  $f$  is the inclusion map. By Proposition 10.14 we have  $\bar{C}_f \simeq Y/X$ , so the above sequence can be written as

$$[X, Z]_* \xleftarrow{f^*} [Y, Z]_* \xleftarrow{q^*} [Y/X, Z]_* \xleftarrow{g^*} [X, \Omega Z]_* \xleftarrow{f^*} [Y, \Omega Z]_* \xleftarrow{q^*} [Y/X, \Omega Z]_* \xleftarrow{g^*} [X, \Omega^2 Z]_* \leftarrow \dots$$

where  $q: Y \rightarrow Y/X$  is the quotient map.